

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
	Abnormality in gait & mobility	781.2	R26.0, R26.1, R26.81, R26.89, R26.9	R26.9		See Category R26 in Tabular List for more detailed diagnosis and code
	Achondroplasia	756.4	Q77.0, Q77.1, Q77.4, Q77.5, Q77.7, Q77.8, Q77.9, Q78.4	Q77.4		
	Amniotic band syndrome	762.8	P02.8	P02.8	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later	
23	Angelman syndrome	759.89	E78.71, E78.72, Q87.2, Q87.3, Q87.5, Q87.81, Q87.89, Q89.8	Q93.5		
	Arthrogryposis	728.3	M62.3, M62.89	Q74.3		
16	Autism/Autistic disorder (childhood, infantile)	299.00	F84.0, F99	F84.0	<b>Use additional code</b> to identify any associated medical condition and intellectual disabilities	
12	Behavior concerns, NOS	313.9 or 312.9	F91.8, F91.2, F91.0, F91.1, F91.9, F93.8, F93.9, F94.8, F98.8, F98.9	F91.9		
	Bronchopulmonary dysplasia	770.7	P27.0, P27.1, P27.8, P27.9	P27.1	<b>Excludes1:</b> respiratory distress of newborn (P22.0-P22.9)	This condition originates in the neonatal period
	Cerebral palsy, infantile, diplegic	343.0	G80.1, G96.8	G80.1		
	Cerebral palsy, infantile, hemiplegic	343.1	G80.2	G80.2		
	Cerebral palsy, infantile, quadriplegic	343.2	G80.0	G80.0		

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
	Cerebral palsy, infantile, unspecified	343.9	G80.0, G80.8, G80.4, G80.9	G80.9		Consult Alphabetic Index and Tabular List for more specific types of Cerebral Palsy
	Child Abuse/Neglect					REFER TO TAB "Child Abuse_Neglect"
	Choanal atresia	748.0	Q89.8, Q30.0	Q30.0		
	Chromosomal anomaly, other	758.9	Q99.9	Q99.8		
	Cleft lip, unspecified	749.10	Q36.0, Q36.1, Q36.9, Q38.0	Q36.9		See Category Q36 in Tabular List for more detailed diagnosis and code (includes laterality)
	Cleft lip and/or palate, history of (corrected)			Z87.730	Excludes1: congenital malformations that have been partially corrected or repaired but which still require medical treatment - code to condition	
	Cleft palate, unspecified	749.00	Q38.5, Q35.1, Q35.3, Q35.5, Q35.9, Q37.0, Q37.1, Q37.2, Q37.3, Q37.4, Q37.5, Q37.8, Q37.9	Q35.9		See Category Q35 in Tabular List for more detailed diagnosis and code
	Cleft palate, unspecified, with bilateral cleft lip	749.23, 749.24	Q35.9, Q37.0, Q37.2, Q37.4, Q37.8, Q38.5, Q38.0, Q35.1, Q35.3, Q35.5, Q36.0	Q37.8		See Category Q37 in Tabular List for more detailed diagnosis and code
	Cleft palate, unspecified, with unilateral cleft lip or cleft lip NOS	749.20	Q35.9, Q37.8, Q37.9, Q36.0	Q37.9		See Category Q37 in Tabular List for more detailed diagnosis and code

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
	Club Foot (congenital)	754.51	Q66.0	Q66.89		Consult Alphabetic Index and Tabular List for more specific types of congenital deformities of feet
	Clonus (Abnormal Reflex)	781.0	R25.0, R25.1, R25.2, R25.3, R25.8, R25.9	R29.2		
	Congenital cytomegalovirus infection (CMV)	771.1	P35.1	P35.1	<b>Includes:</b> infections acquired in utero or during birth	
	Coarctation of the aorta	747.10	Q24.8, Q20.8, Q24.4, Q25.1	Q25.1		
	Coloboma	743.49	Q13.89, Q13.9	Q13.0		See Category Q13 in Tabular List for more detailed diagnosis and code
	Congenital anomaly, unspecified	759.9	Q89.9	Q89.8	<b>Use additional code(s)</b> to identify all associated manifestations	
	Congenital cataract	743.30	Q12.0	Q12.0		
	Congenital infection, Herpes			P35.2	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later	This item is under investigation. Do not use at present time. <b>Question for DHHS Privacy Officer</b>
	Congenital infection, HIV infection status (asymptomatic)			P35.8, Z21	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later	This item is under investigation. Do not use at present time. <b>Question for DHHS Privacy Officer</b>

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
	Congenital infection, HIV infection status (symptomatic, AIDS or ARC)			P35.8, B20	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later	This item is under investigation. Do not use at present time. <b>Question for DHHS Privacy Officer</b>
	Congenital infection, syphilis NOS			A50.2		See Category A50 in Tabular List for more detailed diagnosis and code <b>This item is under investigation. Do not use at present time. Question for DHHS Privacy Officer</b>
	Congenital infection, toxoplasmosis			P37.1		
	Congenital malformations of lung	770.2	NCTracks Crosswalk yields no results	Q33.8		
23	Congenital malformation syndromes	759.89	E78.71, E78.72, Q87.2, Q87.3, Q87.5, Q87.81, Q87.89, Q89.8	See Category Q87 and Q89 in Tabular List	<b>Use additional code(s)</b> to identify all associated manifestations.	
	Corpus Callosum (absent or hypoplastic)	742.2	Q04.0, Q04.1, Q04.2, Q04.3	Q04.0		
	Cortical blindness, unspecified side of brain	377.75	H47.611, H47.612, H47.619	H47.619	<b>Code also</b> underlying condition.	See Subcategory H47.61- for laterality when known
	Cystic fibrosis	277.02	E84.0	E84.9		
24	Dandy-Walker syndrome	742.3	Q03.0, Q03.1, Q03.8, Q03.9	Q03.1		

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
4	Delayed milestones (late walker; late talker)	783.42	R62.0	R62.0		Utilize this diagnosis from time of referral to time of testing. <b>MEDICAID OK with code</b>
1	Developmental delay	315.8	F88	F88		F88 is "Other disorders of psychological development; Developmental agnosia". In the Index, if you look up "Delay, development, global" it sends you to F88. If the CDSAs are unable to be more specific as to the type of delay (Refer to F80 and F82 for more specific speech and motor developmental delay codes), will Medicaid accept F88? <b>MEDICAID OK with code</b>
3	Developmental disorder, motor functions	315.5 or 315.4	F82	F82		
	Developmental disorder, speech and language					Go to "Speech-Language Disorders" below
5	Diagnosis deferred; Undiagnosed disease	799.9	R69, R99	R62.0		According to CDSA staff, Medicaid told them to use ICD-9-CM code 799.9 up to the time for Evaluation. This code translates to R69, "Illness, unspecified; Unknown and unspecified cases of morbidity". R69 is related to morbidity and morbidity does not apply. May CDSAs use R62.0 - Delayed milestones in childhood? <b>MEDICAID OK with code</b>

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
7	Diagnosis, None (No problem found)	V71.9 or V79.3	Z04.9 or Z13.4	TBD - Awaiting test results. Possibilities: Z00.12, Z03.89, Z13.4 or Z71.1		Ideal code would be Z00.12 which includes: Encounter for development testing of infant or child. Excellent description of CDSA initial visits & 6th digit added to clarify with or without abnormal findings; however, use of this code is a billing issue since primary care physicians use this for well visits and there is a limitation on the # of well visits during a year. Other possibilities: Z03.89 – Encounter for observation for other suspected diseases and conditions ruled out; Z71.1 – Person with feared health complaint in whom no diagnosis is made; Z13.4 - Encounter for screening for certain developmental disorders in childhood (says screening and CDSAs do not do screenings (e.g., screening mammogram in asymptomatic person). <b>After discussion with MEDICAID, will include in Testing with NCTracks</b>
	Diaphragmatic hernia (congenital)	756.6	Q79.0, Q79.1	Q79.0	<b>Excludes1:</b> congenital hiatal hernia (Q40.1)	
	DiGeorge's syndrome	279.11	D82.1	D82.1		
9	Down syndrome, unspecified (Trisomy 21 NOS)	758.0	Q90.0, Q90.1, Q90.2, Q90.9	Q90.9	<b>Use additional code(s)</b> to identify any associated physical conditions and degree of intellectual disabilities (F70-F79)	Refer to Category Q90 for more specific diagnosis whenever more specific information is available.
	Dysphagia, unspecified	787.20	R13.0, R13.10	R13.10	<b>Code first</b> , if applicable, dysphagia following cerebrovascular disease (I69. with final characters -91)	Refer to Category R13 for more specific diagnosis whenever more specific information related to the phase is available.

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
	Encephalocele	742.0	Q01.0, Q01.1, Q01.2, Q01.8, Q01.9	Q01.9		
	Examination of ears and hearing	V72.19	Z01.10 or Z01.118	TBD		See "Hearing_Ears" tab
	Epilepsy, NOS	345.0- or 345.9-	G40.309, G40.311, G40.901, G40.909, G40.911, G40.919	G40.909		Refer to Category G40 for more specific diagnosis whenever more specific information related to type of epilepsy is available.
	Erb's Palsy			P14.0		NOTE: Documentation should specify birth injury
6	Expressive language disorder	315.31	F80.1, F80.9, F80.2, F80.4	F80.1		
	Failure to thrive	282.0	D58.0	R62.51		(Failure to gain weight - not newborn)
	Feeding problems of newborn					
	Bilious vomiting of newborn	779.32	P92.01	P92.01		
	Other vomiting of newborn	779.33	P92.09	P92.09		
	Regurgitation and rumination of newborn	779.33	P92.09	P92.1		
	Slow feeding of newborn	779.31	P92.1, P92.2, P92.3, P92.4, P92.5, P92.8, P92.9	P92.2		
	Underfeeding of newborn	779.31	P92.1, P92.2, P92.3, P92.4, P92.5, P92.8, P92.9	P92.3	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs	

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
	Overfeeding of newborn	779.31	P92.1, P92.2, P92.3, P92.4, P92.5, P92.8, P92.9	P92.4	Initially, even if morbidity occurs later <b>Excludes1:</b> feeding problems in child over 28 days old	
	Neonatal difficulty in feeding at breast	779.31	P92.1, P92.2, P92.3, P92.4, P92.5, P92.8, P92.9	P92.5		
	Failure to thrive in newborn	779.34	P92.6	P92.6		
	Other feeding problems of newborn	779.31	P92.1, P92.2, P92.3, P92.4, P92.5, P92.8, P92.9	P92.8		
11	Feeding difficulties/problem (infant - not newborn) (e.g., picky eater, sensory)	783.3	R63.3	R63.3	<b>Excludes1:</b> feeding problems of newborn (P92.-); infant feeding disorder of nonorganic origin (F50.8)	
	Fetal alcohol syndrome	760.71	P04.3, Q86.0	Q86.0		
	Fragile X	759.83	Q99.2	Q99.2		
	Gait abnormality	781.2	R26.0, R26.1, R26.81, R26.89, R26.9	R26.9		See Category R26 in Tabular List for more detailed diagnosis and code
	Gastroschisis	756.73	Q79.3	Q79.3		



### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
17	Hearing impairment /loss (unspecified)	389.9	H91.92, H90.0, H90.2, H90.3, H90.5, H91.01, H91.02, H91.03, H91.09, H91.20, H91.21, H91.22, H91.8x1, H91.8x2, H91.8x3, H91.8x9, H91.90, H91.91, H91.93	TBD		Refer to Category H90 (see "Hearing_Ears" tab) for type of hearing loss (i.e., conductive, sensorineural). Refer to Categories H91 - H93 in code book for other types of hearing loss. Need to specify laterality.
	Hip dysplasia (congenital) without dislocation	755.8	Q74.8	Q65.89		Refer to Category Q65 in code book for more specific types of congenital hip deformities including specifying laterality.
	Hip dysplasia (congenital) with dislocation	754.30	Q65.00, Q65.01, Q65.02, Q65.2	Q65.2		Refer to Category Q65 in code book for more specific types of congenital hip deformities including specifying laterality.
	Hirschsprung's disease (congenital megacolon)	751.3	Q43.1, Q43.2	Q43.1		
	Hydrocephalus, unspecified	331.3	G91.0	G91.9		Refer to Category G91 in code book for more specific types of hydrocephalus; If the client has a shunt, use Status code Z98.2
	Hyperkinesia	314.1	F90.8	R46.3		
	Hypoplastic left heart syndrome	746.7	Q21.0, Q21.4, Q24.8, Q20.8, Q23.4	Q23.4		
8	Hypotonia	781.3	R27.0, R27.8, R27.9	P94.2		

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
	Hypoxic ischemic encephalopathy (HIE), unspecified (newborn)	768.70	P91.60	P91.60	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later	
	Hypoxic ischemic encephalopathy (not newborn)			G93.1		
18	Hypertonia	728.9	M62.9 (NCTracks has a large list including this code)	P94.1	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later	
	Increased Activity	314.1 or 314.9	F90.8 or F90.9	R46.3		
	Infantile spasms, <b>not intractable</b> , without status epilepticus	345.6-	G40.401, G40.409, G40.411, G40.419	G40.822		
	Infantile spasms, <b>intractable</b> , without status epilepticus	345.6-	G40.401, G40.409, G40.411, G40.419	G40.824		
	Infarct/stroke (thrombotic inutero)	434.01	I63.30 (NCTracks has a large list including this code but does not include P96.89)	P96.89 & I63.30	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later <b>Use additional code</b> to specify condition.	
	Intraventricular hemorrhage, unspecified	772.1-	P52.3, P52.0, P52.1, P52.21, P52.22	TBD	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later	Refer to codes P52.0 through P52.22 in code book to determine most accurate code

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
	Intraventricular hemorrhage grade III	772.13	P52.21	P52.21	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later	<b>MEDICAID QUESTION:</b> According to the CDSAs, this condition is currently reimbursable by Medicaid till 1 year of age but coding guidelines allow usage beyond that. Will Medicaid change this rule to coincide with the coding guidelines? The same question applies for prematurity codes. In interim, use up to Age 1.
	Intraventricular hemorrhage grade IV	772.14	P52.22	P52.22	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later	<b>MEDICAID QUESTION:</b> According to the CDSAs, this condition is currently reimbursable by Medicaid till 1 year of age but coding guidelines allow usage beyond that. Will Medicaid change this rule to coincide with the coding guidelines? The same question applies for prematurity codes. In interim, use up to Age 1.
20	Intrauterine growth restriction / retardation (IUGR)	764.90	P05.9	P05.9	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later	REFER TO TAB "Slow Fetal Growth_Malnutrition"
	Lack of Coordination	781.3	R27.0, R27.8, R27.9	R27.8		
19	Low-birth weight					REFER TO TAB "Short Gestation_Low Birth Weight"

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
	Macrocephaly (congenital)	756.0	Q75.0, Q75.1, Q75.2, Q75.3, Q75.4, Q75.5, Q75.8, Q75.9, Q87.0	Q75.3		
25	Malrotation (intestine)	754.1	Q68.0	Q43.3		
	Meningitis (bacterial), History of	320.9	G00.9, G04.2	Z86.61		
	Metabolic disorder	277.9	NCTracks Crosswalk yields no results	TBD		Consult with Medical Staff - refer to Categories E70-E88 in code book to determine most accurate code
	Microcephaly (congenital)	742.1	Q02	Q02		
2	Mixed receptive-expressive language disorder	315.31 or 315.32	F80.1, F80.2, F80.9, F80.4, H93.25	F80.2		
	Necrotizing colitis	777.50	P77.9	P77.9	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later	
	Neonatal seizures	779.0	P90	P90		
	Nystagmus, unspecified	379.50	H55.00	H55.00		
	Optic nerve hypoplasia, <b>right</b> eye	377.43	H47.031, H47.032, H47.033, H47.039	H47.031		
	Optic nerve hypoplasia, <b>left</b> eye	377.43	H47.031, H47.032, H47.033, H47.039	H47.032		
	Optic nerve hypoplasia, <b>bilateral</b>	377.43	H47.031, H47.032, H47.033, H47.039	H47.033		

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
	Other perinatal conditions, specified	779.8-	P29.12, P29.11, P96.82, P96.83, P29.81, P29.0, P29.89, P94.1, P94.2, P96.89	P96.89	Includes: conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later <b>Use additional code</b> to specify condition	What should be used if a client has met the criteria for enrollment and they are entitled to Early Intervention services up to age 3 but the perinatal condition no longer exists? Child may be receiving case management services only. <b>Question for Medicaid</b>
	Periventricular leukomalacia (PVL)	779.7	P91.2	P91.2	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later	
	Pervasive developmental disorders (PDD)	299.80	F84.2, F84.5, F84.8, F84.9, F81.9	F84.9		
	Pierre Robin Syndrome	756.0	Q75.0, Q75.1, Q75.2, Q75.3, Q75.4, Q75.5, Q75.8, Q75.9, Q87.0	Q87.0		
	Plagiocephaly	754.0	Q67.0, Q67.1, Q67.2, Q67.3, Q67.4	M95.2		Acquired
	Prader Willi	759.81	Q87.1	Q87.1		
19	Prematurity					REFER TO TAB "Short Gestation_Low Birth Weight"
	Pulmonary atresia (artery)	747.31	Q25.5, Q25.71, Q89.8, Q25.79	Q25.5		
	Pulmonary atresia (valve)	747.31	Q25.5, Q25.71, Q89.8, Q25.79	Q22.0		
	Schizencephaly	742.4	Q00.0, Q04.4, Q04.5, Q04.6, Q04.8, Q89.8	Q04.6		

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
10	Screening for Developmental problem	V79.3	Z13.4	DO NOT USE		Screening is the testing for disease or disease precursors in asymptomatic individuals so that early detection and treatment can be provided for those who test positive for the disease. CDSAs do not screen.
	Seizure disorder NOS	345.9-	G40.901, G40.909, G40.911, G40.919	G40.909		
21	Sensory disturbances	782.0	R20.0, R20.1, R20.2, R20.3, R20.8, R20.9	TBD		<b>QUESTION FOR Sherry Franklin-</b> Determination needed on this. Should CDSAs be seeing children with sensory disturbances?
22	Septo-optic dysplasia	742.2	Q04.0, Q04.1, Q04.2, Q04.3	Q04.4		
	Shaken infant syndrome	995.55	T74.4xxA	T74.4xxS	A=initial encounter; D=subsequent encounter; S=sequela	Code assumes all CDSA clients will be sequela. Also see tab "Child Abuse_Neglect"
	Short attention span			R45.87		
19	Short gestation					REFER TO TAB "Short Gestation_Low Birth Weight"
	Short-gut (bowel) syndrome	579.3	K91.2	K91.2		
	Short stature	783.43	R62.52	R62.52		
	Sleep disorder (nonorganic)	307.40	F51.9	F51.9		
14	Speech-language therapy					REFER to Tab "Hearing_Ears"
	Speech-language disorders					
	Developmental disorder, speech and language - Phonological disorder			F80.0		
	Developmental disorder, speech and language - Expressive language disorder			F80.1		

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
2	Developmental disorder, speech and language - Mixed receptive-expressive language disorder			F80.2		
	Developmental disorder, speech and language - Speech and language developmental delay due to hearing loss			F80.4	Code also type of hearing loss (H90.- , H91.-)	
	Developmental disorder, speech and language - Childhood onset fluency disorder (cluttering, stuttering)			F80.81		
	Developmental disorder, speech and language - Other			F80.89		
	Developmental disorder, speech and language - Unspecified			F80.9		
	Spina bifida (unspecified) <b>without</b> hydrocephalus	741.90	Q05.9, Q05.8, Q07.00	Q05.9		Refer to Category Q05 in code book for more specific types of hydrocephalus
	Spina bifida (unspecified) <b>with</b> hydrocephalus	741.00	Q05.1, Q05.2, Q05.3, Q05.4, Q05.9, Q07.01, Q07.02, Q07.03	Q05.4		Refer to Category Q05 in code book for more specific types of hydrocephalus
	Spinal muscular atrophy, infantile, type 1 (Werdnig-Hoffman)	335.10	G12.9	G12.0		
	Spinal muscular atrophy, other inherited	335.10	G12.9	G12.1		

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
	Subdural hematoma, traumatic, NOS	852.2-	S06.5x--	S06.5x9S	Use secondary code(s) from Chapter 20, External causes of morbidity, to indicate cause of injury.	
	Stroke/infarct (thrombotic)	434.01	I69.30	P96.89 & I69.30	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later <b>Use additional code</b> to specify condition.	I69.30 - Unspecified sequelae of cerebral infarction may be more appropriate for CDSA clients since active treatment for the stroke has probably concluded when CDSAs see the client.
	Tethered spinal cord (acquired)			TBD		<b>Look at Category G95 - are any of these codes appropriate?</b>
	Tethered spinal cord (congenital)	742.59	Q06.0, Q06.1, Q06.2, Q06.3, Q06.4, Q06.8, Q06.9, Q05.9, Q07.00	Q06.8		
	Tetralogy of Fallot	745.2	Q21.0, Q21.1, Q21.2, Q21.3, Q21.4, Q21.8, Q21.9	Q21.3		
	Torticollis (acquired)			M43.6	<b>Excludes1:</b> congenital (sternomastoid) torticollis (Q68.0); spasmodic torticollis (G24.3); torticollis due to birth injury (P15.2)	
25	Torticollis (congenital)	754.1	Q68.0	Q68.0	<b>Excludes1:</b> reduction defects of limb(s) (Q71-Q73) <b>Excludes2:</b> congenital myotonic chondrodystrophy (G71.13)	
	Tracheoesophageal fistula (congenital)	750.3	Q39.0, Q39.1, Q39.2, Q39.3, Q39.4	Q39.2		



### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
	Trisomy 13, unspecified	758.1	Q91.4, Q91.5, Q91.6, Q91.7	Q91.7		Refer to Category Q91 for more specific diagnosis whenever more specific information is available.
	Turner's syndrome	758.6	Q96.0, Q96.1, Q96.2, Q96.3, Q96.4, Q96.8, Q96.9	Q96.9	<b>Excludes1:</b> Noonan syndrome (Q87.1)	Refer to Category Q96 for more specific diagnosis whenever more specific information is available.
	Underimmunization status			Z28.3	<b>Includes:</b> Delinquent immunization status; Lapsed immunization schedule status	
23	Vacterl/Vater association/syndrome	759.89	E78.71, E78.72, Q87.2, Q87.3, Q87.5, Q87.81, Q87.89, Q89.8	Q87.2	identify all associated manifestations. <b>Includes:</b> Holt-Oram syndrome; Klippel-Trenaunay-Weber syndrome; Nail patella syndrome; Rubinstein-Taybi syndrome; Sirenomelia syndrome; Thrombocytopenia with absent radius [TAR] syndrome; VATER syndrome	
	Ventricular septal defect (congenital)	745.4	Q21.0, Q21.8	Q21.0		
	Vision loss, unspecified	369.9	H54.7	H54.7		Refer to Category H54 for more specific diagnosis whenever more specific information is available.
	Vocal cord paresis	478.31	J38.01	J38.00	<b>Use additional code</b> to identify: exposure to environmental tobacco smoke (Z77.22) exposure to tobacco smoke in the perinatal period (P96.81)	Refer to Category J38 for more specific diagnosis whenever more specific information is available.

### CDSA Common Dx Reference List v3.3

NOTE: All diagnosis codes should be validated using the Tabular List

Frequency of Dx in CDSAs	Diagnostic Description	ICD-9-CM Code	NCTracks Crosswalk Codes	ICD-10-CM Code	ICD-10-CM Tabular Instructions	Questions/Comments
	(Neonatal) Withdrawal symptoms from maternal use of drugs of addiction			P96.1	<b>Includes:</b> conditions that have their origin in the fetal or perinatal period (before birth through the first 28 days after birth) even if morbidity occurs later <b>Excludes1:</b> reactions and intoxications from maternal opiates and tranquilizers administered during labor and delivery (P04.0)	This item is under investigation. Do not use at present time. <b>Question for DHHS Privacy Officer</b>